
Dr. Weiss's Opinions

- Adverse event reports cannot reliably be used to establish a causal relationship between gabapentin and suicide
- The FDA's AERS database does not support the finding of a signal for completed suicide or suicide attempt
- Dr. Blume's methodology for reviewing AERS data is flawed and not generally accepted
- The information in the AERS database for gabapentin and suicide is unreliable and therefore uninterpretable


Sample MedWatch Form



Submission of a report does not constitute an admission that medical personnel or the product caused or contributed to the event.

Case 3:05-cv-00444 Document 181-2 Filed 04/27/10 Page 2 of 15 PageID #: 4372

Data Mining Cannot Establish Causation




March 2005

Data mining is not a tool for establishing causal attributions between products and adverse events.

Source: March 2005 FDA Guidance for Industry: Good Pharmacovigilance Practices and Pharmacoepidemiologic Assessment, Pg. 8

Clinical Judgment Necessary For Review of Data Mining




March 2005

Proper interpretation also requires clinical judgment before one even considers there to be a signal.


Source: Brian L. Strom, M.D., MPH, Reply to Letter to the Editor, 293 JAMA (2005), Pg. 1325

Case 3:05-cv-00444 Document 181-2 Filed 04/27/10 Page 4 of 15 PageID #: 4374


Gabapentin Spontaneous Reports Over Time



Neurontin/Gabapentin Prescriptions Over Time




Gabapentin Reports as a Percentage of Total Prescriptions




Case 3:05-cv-00444 Document 181-2 Filed 04/27/10 Page 7 of 15 Page Duttice 4 Weiss Expert Report, Figure 3


Gabapentin Reports Decrease As Prescriptions Increase




Proportion of Reports In Which Gabapentin 'Was Suspect'



Gabapentin Reports by Report Type – By Year




Gabapentin Reports by Report Type – By Quarter




Case 3:05-cv-00444 Document 181-2 Filed 04/27/10 Page 11 of 15 Pages Drte: West Expert Report, Figure 7

PRR for Gabapentin: Completed Suicide And Suicide Attempt



FDA: Controlled Trials Are Only Way to Assess Whether Neurontin Is Associated With Increased Risk of Suicide





April 12, 2005 Letter From FDA to Plaintiff's Lawyers

With the letter, you provided 258 MedWatch forms of patients whom you state committed suicide while being treated with Neurontin, and you make several allegations about FDA's handling of this matter.


* * *

We noted at that time that these illnesses are well-known to be associated with an increased risk of suicide compared to the general population. Further, in the absence of an appropriate control group, it will be difficult, if not impossible, to assess the role of any other factors that might explain these events, such as concomitant medications.

Source: April 12, 2005 Letter from FDA

Case 3:05-cv-00444 Document 181-2 Filed 04/27/10 Page 13 of 15 Page D #. 4383

Gabapentin vs. FDA AED Drugs



Completed Suicide Reports



